



The Open Nursing Journal

Content list available at: <https://opennursingjournal.com>



RESEARCH ARTICLE

Comparison of the Effect of Aerobic and Resistance Training on Cardiovascular Risk Factors in Parkinson's Women

Hamideh Yaghoobi¹, Tahere Sarbooz Hossein-Abadi¹, Sahar Mohammadi², Samaneh Sardashti^{1,*} and Seddigheh Abbaspour¹

¹Department of Nursing, School of Nursing and Midwifery, Torbat Heydariyeh University of Medical Sciences, Torbat Heydariyeh, Iran

²Department of Physical Education and Sport Sciences, Islamic Azad University, Najafabad, Iran

Abstract:

Background:

People with low-mobility are at increased risk for cardiovascular disease. Patients with Parkinson's Disease (PD) are less interested in participating in exercise activities due to the nature of the disease. Regular exercise and physical activity can play protective roles against heart disease by reducing risk factors. This study aimed to compare the effects of aerobic and resistance training on cardiovascular risk factors in women with PD.

Materials and Methods:

This quasi-experimental study was carried out on 45 women with PD who were randomly assigned to one of three groups: aerobic, resistance groups, and control group. Before and after training sessions, variables of weight, Body Mass Index (BMI), heart rate at rest state, maximum oxygen consumption, blood pressure, and lipid profiles of participants were measured. Data were analyzed using SPSS software version 21 and Paired Sample t-test, ANOVA and Bonferroni post-hoc test. A significant level of $P \leq 0.05$ was considered.

Results:

Demographics of physiological characteristics of subjects was congruent in all three groups in the baseline. After aerobic and resistance training, levels of triglyceride, LDL, total cholesterol, fat percentage, resting heart rate, weight, BMI and systolic blood pressure were significantly decreased ($p < 0.05$). This change was more pronounced in the aerobic group; in contrast, variables of maximum oxygen consumption and HDL were meaningfully increased in participants of both experimental groups ($p < 0.05$).

Conclusion:

The results showed that both types of aerobic and resistance training can reduce the risk factors of cardiovascular disease in women with PD, while the effect of aerobic training on reducing risk factors was more than resistance training.

Keywords: Parkinson disease, Resistance training, Aerobic exercise, Cardiovascular diseases, Risk factors, Alzheimer's disease.

Article History

Received: July 06, 2019

Revised: October 15, 2019

Accepted: October 14, 2019

1. INTRODUCTION

Parkinson's Disease (PD) is in the group of degenerative and progressive diseases, and with slow movement in the Central Nervous System (CNS). In the current century, it is the second common disease of the CNS system after Alzheimer's disease, and is one of the most common causes of disability in old age [1]. PD affects men more than women, and its symptoms are usually manifested in the fifth decade of life. In the United States, this disease is caused by neurodegenerative

disease, and annually 60,000 new cases are reported [2, 3]. In Iran, due to the growth of the elderly population relative to the total population, the disease is expected to increase [4].

The two-rings reducing and defective of PD including aging and inactivity may lead to increased complications of the disease and secondary problems for patients [5]. One of these complications is cardiovascular disease, which has increased in recent decades and is considered as one of the main causes of mortality in the world [6]. About forty-six percent of deaths result from chronic diseases, especially in old age [7].

With respect to numerous scientific evidences, inactivity,

* Address correspondence to this author at the Department of Nursing, School of Nursing and Midwifery, Torbat Heydariyeh University of Medical Sciences, Iran; E-mail: sardashtis1@thums.as.ir

especially in the new age has become a problem that it has been included in the list of cardiovascular risk factors [8]. Previous evidences show that low-mobility life increases the risk of coronary heart disease therefore, sedentary people are twice more likely to suffer from the risk of coronary artery disease than the active people [9]. Several studies have shown that inactivity increases cardiovascular risk factors in people, disrupts the quality of life, and eventually leads to cardiovascular diseases. In a study by Mora *et al.*, the correlations between physical activity and Body Mass Index (BMI) with cardiovascular indices were measured, they concluded lower physical activity and an increase in BMI independently associated with increased Total Cholesterol (TC), Triglyceride (TG) and Low-density lipoprotein cholesterol (LDL-c) [10]. The findings of the Gidding study showed that physical inactivity can cause an increase in cardiovascular risk factors and insulin resistance [11].

In order to cope with this huge and growing problem of human societies, it seems that safer and more appropriate solutions should be discovered, in addition to medication. One of these methods is exercise treatments. Contrary to the previous literature which showed that exercise caused physical stress and negatively affected the heart, recent research findings suggest that physical activity is safe and beneficial even for people with functional impairment and patients with heart failure [12]. Additionally, regular exercise can lead to increased independence for daily activities, prevention of functional disability and dependency on specific situations. Physical activity can be used as a valuable tool in the effective treatment of heart disease symptoms, and improve the functional capacity and the life quality of patients [13, 14].

Because of the nature of the disease, it seems that patients with PD are less interested in participating in exercise activities [15]. One way to help these patients is through exercise. Systematic participation in a physical activity program can help individuals to not only maintain their motor resources but also gain the ability to perform activities in their daily life. On the other hand, regarding the growth of the elderly population in Iran and the possible increase in the number of patients with PD, this study aimed to compare the effect of aerobic and resistance training on cardiovascular risk factors in women with PD.

2. MATERIALS AND METHODS

2.1. Subjects

The present study was a semi-experimental study with pre-test-post-test design and control group. In this study, forty-five women with PD referring to the Neurology clinic of two Hospital Torbat Heydariyeh voluntarily participated. The participants were randomly assigned three groups: aerobic training, resistance training, and control group (fifteen people in each group). Before the training, all subjects stated their written consent in a satisfactory manner. All procedures performed in this study were in accordance with the ethical committee of Torbat Heydariyeh University of Medical Sciences and with the principles of the declaration of Helsinki (1964) and its later amendments or comparable ethical standards.

The criteria for entering the study were to have PD for more than a year, a sedentary lifestyle, and the use of the same drugs at the same dose according to the neurologist. The exclusion criteria included BMI ≥ 43 (based on the highest BMI belonging to people capable of exercising), age > 70 years, history of brain and cardiovascular diseases, as well as the history of musculoskeletal problems limiting physical activity.

2.2. Method of Exercising

2.2.1. Implementation of Resistance Training

The resistance group training program included a series of resistance exercises that were conducted lasted for 8 weeks, 3 sessions per week, and each session for 60 minutes (Table 1). In the first session, the basic exercises were explained for subjects, which the principles of course were respected at all meetings. At the beginning of each session, the training sessions included warming (about 10 minutes), the continuing session was followed by performing resistance exercises (about 40 minutes), and at the end of the session, refreshment was followed by a return to the initial state (about 10 minutes).

To familiarize and train the correct implementation of resistance exercises, all subjects fulfilled several sub-maximal repetitions for each movement in the first session. In the next session, to calculate the load intensity of exercise for each subject, a single maximal repeat test was performed for every 6 movements of the exercise program, these movements included: chest press, dumbbell press, triceps pushdown, abdominal crunch, leg press, thigh front, and thigh behind. It is necessary to note that the used weight was not maximized during repetitions. And according to the ability of the subjects, by observing the selected safety points, the Brzycki equation was used for the calculation of one repetition maximum (1RM) [1, 16].

These exercises were done in a circle with 40-60% maximum repetition. Since the participants were not familiar with resistance training and had weak ability, the exercises were started from a very low level and were gradually progressed. In order to comply with the principle of overload, the movement repetition in each session was increased as compared to the previous meeting.

Table 1. Resistance group training programs.

Movements	Set	Repeat	Intensity (One Repetition Maximum)	Number of Exercise Times
First and second weeks	3	12-15	40-45	2-4
Third and fourth weeks	3	10-12	45-50	2-4
Fifth and Sixth weeks	3	8-10	50-55	2-4
Seventh and eighth week	3	5-8	55-60	2-4

2.2.2. Implementation of Aerobic Exercise

For aerobic exercises, the intensity of activity was Heart Rate Reserve (HRR) 50% and its duration was 20 minutes, with 12 minutes on the treadmill and 8 minutes on the Bike Ergometer. The activity on the Bike Ergometer was determined

by Revolutions per Minute (RPM). RPM is one of the angular velocity measuring units that express the number of complete turning cycles in a single minute around a constant axis [17]. The subjects initially started with 30 to 40 RPM and each week gradually increased the intensity of their activity. Also, the activity time was added 2 minutes every week, and in the eighth week, it reached 70% HRR and 30 minutes of activity.

The subjects of the control group underwent usual drug treatments during the 8-week training program and only performed routine activities. However, experimental subjects in addition to drug therapy fulfilled 3 sessions of resistance and aerobic exercises for 60 minutes during the 8-week. It should be noted that exercises by adhering to the scientific principles of research were carried out under the supervision of the trainer. The studied physical and physiological variables included weight, BMI, Resting Heart Rate (RHR), maximum oxygen consumption, systolic and diastolic blood pressure, and risk factors of cardiovascular disease such as LDL-C, High-Density Lipoprotein Cholesterol (HDL-C), TC, and TG, their changes were measured and recorded by applying independent variables on subjects for 8 weeks.

2.2.3. Measurement Tools

The measurement tools in this study included strip meter, stadiometer model seca 208 with 1% cm accuracy, seca portable scale with a precision of 1% kg, treadmill test on treadmill device model technology made in Italy for measurements of aerobic capacity, Barometer model omron made in Vietnam was used for measuring blood pressure and heart rate monitor model H6 made in Finland was used to measure the heart rate, and the kit of Pars test Com. was used to measure blood lipids.

To determine blood lipids after 8-10 hours of fasting at 8 am (48 hours before physical activity and 48 hours after the end of exercise), blood samples of individuals were taken in an open arch for determining the level of cholesterol, LDL, HDL. The measurements of cholesterol and TG were done using the colorimetry method, and the variables of LDL and HDL were measured using colorimetry-direct method, by the Hitachi 912 device Roche Com. made in Germany.

2.2.4. Statistical Analysis

The quantitative data including age, height, weight, and *etc.* were expressed as mean and standard deviation ($M \pm SD$). The normality of data was confirmed by Kolmogorov-Smirnov test. Data analysis was performed using SPSS software version 21. The Paired Sample t-test was used to differentiate between two measurements pre-test and post-test in each of the groups. Because the comparison of three groups means that the one-

way analysis of variance (ANOVA) was utilized, and the Bonferroni posthoc test was also used to detect differences between groups in the studied variables. A significant level of $p \leq 0.05$ was considered.

3. RESULTS

Table 2 shows the comparison of demographic variables of subjects including age, height, weight, and BMI before the study. Based on ANOVA, there was no significant difference between the mean of the demographic characteristics of subjects in all the three groups. Table 3 exhibits a comparison of physiological variables and lipid profiles, such as TG, TC, LDL-C and HDL-C between the experimental groups (aerobic and resistance training) and control group before the study. ANOVA Analysis also showed no significant difference between the mean of physiological characteristics and lipid profile of subjects in all three groups ($P > 0.05$). Therefore, all three groups were congruent comparing the mean of intra-group and inter-group differences of the studied variables in the studied groups is presented in Table 4. The results of Paired Sample t-test indicated that variables of weight ($P < 0.001$), BMI ($P = 0.01$), body fat percentage (BFP) ($P = 0.01$), heart rate ($P = 0.01$), maximum oxygen consumption ($P < 0.001$), LDL ($P = 0.01$), HDL ($P < 0.001$), and TC ($P = 0.01$) in patients with PD significantly declined in aerobic group trained for eight weeks. Also, after eight weeks of resistance training compared with the baseline, maximum oxygen consumption ($P = 0.01$) and HDL ($P < 0.04$) significantly diminished in Parkinson's patients. There was no significant difference between the mean of TG in PD patients in studied groups after aerobic and resistance training compared to baseline. This difference was also not significant in the control group ($P > 0.05$).

The results of ANOVA test showed that there were significant differences between variables of weight ($P = 0.007$), BMI ($P = 0.02$), body fat percentage ($P < 0.001$), heart rate ($P < 0.001$) maximum oxygen consumption ($P < 0.001$), HDL ($P < 0.001$), and TC ($P < 0.001$) among studied groups; while no significant difference was observed between LDL and TG variables among studied groups ($P > 0.05$), as shown in Table 4.

A comparison of Bonferroni post-hoc test between studied groups in Parkinson's patients is shown in Table 5. On applying statistical analysis, results revealed that there are significant differences between the mean of heart rate, oxygen consumption, systolic and diastolic blood pressure, fat percentage, HDL, and TC in experimental groups in compared control group (all $P < 0.05$). BMI of aerobic exercise group was significantly different from the control group ($P = 0.02$) and there was also a significant difference between the weight of subjects with aerobic training in comparison to resistance training and control groups ($P < 0.05$).

Table 2. Comparison of the mean and standard deviation of physical characteristics of subjects in studied groups before study.

Variable	Aerobic Exercise	Resistant Exercise	Control Group	F	p-value
Age (years)	49.6±33.49	49.6±20.27	49.7±53.07	0.13	0.94
Height (cm)	161.6±97.9	161.4±12.51	160.5±25.8	1.08	0.36
Weight (kg)	85.7±80.87	85.7±2.89	85.7±60.06	0.04	0.99

(Table 2) contd.....

Variable	Aerobic Exercise	Resistant Exercise	Control Group	F	p-value
BMI (kg/m ²)	29.2±20.34	28.2±25.09	27.1±86.79	0.91	0.45
Fat (%)	31.3±52.28	30.2±64.76	31.3±71.74	0.41	0.75

Table 3. Comparison of the mean and standard deviation of physiological characteristics and lipid profile of subjects in studied groups before.

Variable	Aerobic Exercise	Resistant Exercise	Control Group	F	P-value
Heart rate (bpm)	77.73±8.06	77.73±6.74	79.06±9.54	0.36	0.78
maximal oxygen consumption (ml/kg/min)	19.67±6.47	20.48±5.92	18.96±5.16	0.84	0.48
Triglyceride (mg/dl)	139.6±51.3	142.8±41.56	145.18±44.6	0.25	0.86
Cholesterol (mg/dl)	275.57±28.23	268.57±23.6	276.73±33.4	0.36	0.78
LDL (mg/dl)	128.67±27.69	123.2±24.43	125.2±27.48	0.13	0.95
HDL (mg/dl)	39.2±9.1	38.66±5.99	38.73±7.25	0.18	0.91

Table 4. Intra-group and inter-group differences for comparing the mean of the studied groups.

Variable	Group	Test State		Intra-Group		Between Groups	
		Before	After	t	P-value	F	P-value
Weight (kg)	Aerobic	85.8±7.88	83.83±8	4.612	0.001	5.642	0.007
	Resistant	85.73±7.9	85.49±7.7	0.702	0.49		
	Control	85.6±7.07	85.83±6.5	-0.345	0.74		
BMI (kg/m ²)	Aerobic	29.03±2.35	28.35±2.29	4.437	0.01	4.631	0.02
	Resistant	28.2±2.09	28.18±2.08	0.68	0.51		
	Control	27.87±1.79	27.94±1.61	-0.369	0.72		
Fat (%)	Aerobic	31.52±3.28	28.65±4.31	4.437	0.01	14.966	0.001
	Resistant	30.65±2.77	29.75±3.41	0.68	0.51		
	Control	31.71±3.74	34.69±3.71	-0.369	0.72		
Heart rate (bpm)	Aerobic	78.73±10.63	71.67±10.63	4.437	0.01	14.916	0.001
	Resistant	76.73±6.74	74.33±8.43	1.102	0.29		
	Control	79.06±9.54	86.47±8.73	-3.382	0.004		
maximal oxygen consumption (ml/kg/min)	Aerobic	19.68±3.39	23.68±3.41	-7.407	0.001	35.844	0.001
	Resistant	20.49±3.92	21.64±3.94	-2.921	0.01		
	Control	18.71±5.16	16.34±4.96	3.362	0.05		
LDL (mg/dl)	Aerobic	128.67±27.7	118.33±26.86	4.437	0.01	1.16	0.32
	Resistant	123.2±24.43	117.93±16.99	1.102	0.29		
	Control	125.2±29.49	124.93±22.19	-3.382	0.004		
HDL (mg/dl)	Aerobic	39.2±9.1	42.27±9.32	-5.193	0.001	18.935	0.001
	Resistant	38.67±5.99	42.6±7.07	-2.227	0.04		
	Control	38.73±7.26	33.27±5.78	3.034	0.009		
Cholesterol (mg/dl)	Aerobic	275.57±28.23	250.83±37.22	4.143	0.01	14.252	0.001
	Resistant	268.57±23.61	260.17±29.51	1.102	0.29		
	Control	276.73±33.4	301.63±30.13	-3.205	0.006		
Triglyceride (mg/dl)	Aerobic	139.6±51.31	135.87±41.61	0.823	0.42	1.037	0.36
	Resistant	142.8±41.56	138.93±23.73	0.421	0.68		
	Control	153.2±44.6	155.47±48.27	-1.081	0.3		
Systolic blood pressure (mmHg)	Aerobic	137.5±98.31	115.12±7.1	5.185	0.000	18.319	0.001
	Resistant	144.8±9.4	120.1±5.2	2.153	0.001		
	Control	123.2±10.6	130.2±8.62	3.045	0.001		
Diastolic blood pressure (mmHg)	Aerobic	82.10±4.5	52.3±2.21	4.215	0.001	15.056	0.001
	Resistant	80.7±4.82	61.4±2.56	3.143	0.002		
	Control	70.1±5.05	72.3±6.75	3.012	0.001		

Table 5. Comparing the Bonferroni post hoc test between studied groups in Parkinson's patients.

Variable	Group I	Group J	Mean Difference (I-J)	P-value
Weight (kg)	Aerobic	Resistant	-1.729	0.04
		Control	-2.188	0.008
BMI (kg/m ²)	Aerobic	Control	-0.66	0.02
Fat (%)	Aerobic	Control	-5.896	0.001
	Resistant	Control	-4.159	0.002
Heart rate (bpm)	Aerobic	Control	-14.572	0.001
	Resistant	Control	-10.532	0.001
maximal oxygen consumption (ml/kg/min)	Aerobic	Control	6.514	0.001
	Resistant	Control	3.784	0.001
HDL (mg/dl)	Aerobic	Control	12.703	0.001
	Resistant	Control	9.376	0.001
Cholesterol (mg/dl)	Aerobic	Control	-50.015	0.001
	Resistant	Control	-35.973	0.002
Systolic blood pressure (mmHg)	Aerobic	Control	-10.564	0.001
	Resistant	Control	-4.526	0.002
Diastolic blood pressure (mmHg)	Aerobic	Control	-9.376	0.001
	Resistant	Control	-2.246	0.000

4. DISCUSSION

Since the main emphasis of rehabilitation programs is on exercise and physical activity, the present study was designed for evaluating and comparing the effects of aerobic and resistance training on cardiovascular risk factors in women with Parkinson's. In order to prevent any possible injury during training, all exercises were performed under the supervision of a trainer. The results of this study exhibited that 8 weeks of resistance and aerobic training reduced the levels of LDL, TG, TC, fat percentage, RHR, weight and BMI, and also increased HDL and oxygen consumption, which indicates the importance of exercise and physical activity in protecting the health and treatment of Parkinson's patients. According to the results, aerobic training had a significant effect on reducing cardiovascular risk factors compared with resistance training which is consistent with the findings of Taheri *et al.* [18]. They showed that aerobic training is effective in reducing TC levels. Also, Mogharnasi and Bagheri observed the significant effects of aerobic and resistive exercises on the reduction of blood cholesterol levels [14]. In addition, Soori *et al.* indicated that resistance and endurance training significantly reduced cholesterol and TG levels in the aerobic and resistance groups [19]. The results of a systematic review showed the positive effects of exercise training on reducing glycated hemoglobin and other risk factors, and stated that hybrid exercises are more effective than aerobic or resistance training alone [20]. However, the results of the studies by Nayebifar *et al.* and Hosseini [21, 22], were not consistent with the findings of this study. In their study, there was no relationship between exercise and lipid profile. The probable cause of this inconsistency was related to the severity and the type of exercises. In order to reduce the plasma TG, the duration of exercise should be long, although in some cases, lessening TG is also affected by the severity of exercise. TG reduction is likely due to an increase in lipoprotein lipase activity of muscle capillaries [23]. Furthermore, the effect of exercise on the

concentration of plasma TG depends on concentration prior to the activity. Individuals with low levels of TG did not show significant changes in exercise activity, while plasma TG in subjects with high concentrations significantly reduced [24]. Moderate-intensity exercises are needed to maximize the benefits of improving blood fat levels. On the other hand, LDL levels in the study by Saadatnia *et al.* showed that aerobic and resistance training had decreasing effects on the half-lives of apolipoproteins and lipoproteins [25]. Similar results were found by Baharloo *et al.* which showed that exercise has positive effects in lowering the LDL levels [26].

In this study, a significant correlation was observed between physical activity and HDL levels, which is similar to the results of Askari *et al.* and Maggio *et al.* [27, 28]. They found that exercise can be effective in increasing HDL levels. In addition, Resaland *et al.* [29] showed a significant positive relationship between exercise and HDL levels. All of these studies have been associated with an increase in HDL levels, which is in agreement with the results of this study. However, the results of our study were inconsistent with the findings of Azizi *et al.*, and Rahmani-Nia *et al.* [30, 31]. Azizi *et al.* investigated the relationship between physical activity level and cardiovascular risk factors in male students, and Rahmani-Nia *et al.* studied the effects of exercise on serum leptin and risk factors for coronary heart disease. Regular physical activity can be effective for the cardiovascular system, blood pressure, body fat percentage, and body fitness. Therefore, Abdi *et al.* found that aerobic exercises had effects on body composition and plasma levels of Insulin-like growth factors and their binding proteins, and likewise, BMI, lipid percentage, weight, waist-to-hip ratio after 8 weeks of aerobic training and maximum oxygen consumption in experimental group were significantly increased [32]. Furthermore, Attarzadeh Hoeyini *et al.* observed that aerobic exercises can increase capacity and respiratory coordination, especially in muscles of exhalation by enriching lung's capacity and volume. Exercise can improve

lung function and increase oxygen consumption [33].

Here, heart rate and blood pressure of the control group were significantly higher at the end of the study, which may be due to the possible effects of medications [34, 35]. The results of the study by Najaran and colleagues showed that levodopa may have possible effects to cause or worsen the autonomic nervous system dysfunction in some patients [34]. Also, Kazemi and Rafatpanah observed the paradoxical effects of some drugs on blood pressure in the PD [35].

Based on the results, exercise can be effective in controlling systolic and diastolic blood pressure. In a study conducted by Rahimian *et al.* A 16-session aerobic exercise program with a six-week diet was considered, in which diastolic, systolic, and mean arterial pressures were reduced to 8.5mmHg, 20mmHg, and 12mmHg, respectively. Moreover, the maximum oxygen consumption was improved to 5.7ml/kg/min [36, 37]. In the present study, systolic blood pressure was significantly decreased in both aerobic and resistance groups compared to the control group. Diastolic blood pressure was also changed in aerobic and resistance groups, but this change was not significant. In the control group, diastolic blood pressure was almost constant. Johansen demonstrated that aerobic exercises were effective in reducing and controlling the blood pressure of kidney patients [38]. The results of Smeltzer *et al.* indicated a decrease in systolic blood pressure in hypertensive patients with physical activity [39]. Nonetheless, the results of studies by Myers *et al.* and Rafati *et al.* [40, 41] did not conform with this study. They found that exercise does not have an effect on blood pressure control.

This research was conducted by a small sample size on women referred to the clinic with respect to financial and time resources, which are limitations of the study.

CONCLUSION

The results showed that both types of aerobic and resistance training can reduce the risk factors of cardiovascular disease in women with PD. However, the effect of aerobic training on reducing risk factors was more than resistance training. Therefore, exercise can be effective as a therapeutic option in reducing the risk factors of cardiovascular disease and it is recommended along with medical treatment.

ETHICAL APPROVAL AND CONSENT TO PARTICIPATE

This research was approved by the Ethics Committee at the Torbat Heydariyeh University of Medical Sciences (Code Number IR.THUMS.REC.1397.0021).

HUMAN AND ANIMAL RIGHTS

No animals/humans were used for studies that are the basis of this research.

CONSENT FOR PUBLICATION

Informed consent was obtained from all participants.

AVAILABILITY OF DATA AND MATERIALS

The data that support the findings of this research are

available from corresponding author upon request with permission from Ethics Committee of THUMS.

FUNDING

This work was supported by the the Torbat Heydariyeh University of Medical Sciences (No. A-10-1311-21).

CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

ACKNOWLEDGMENTS

The authors appreciate the “Mahoor Sport Club” and “all the staff of the neurology clinics of the 9-Day hospital and the social security hospital” in Torbat Heydariyeh, which helped us in this research.

REFERENCE

- [1] Mostafavian Z, Gohardehi F, Shakerian S, *et al.* Quality of life in patients with Parkinson's disease: Assessing with the Parkinson's Disease Questionnaire (PDQ-39). *Majallah-i Ulum-i Pizishki-i Razi* 2012; 19(94): 33-9. Available at: <http://jrms.mui.ac.ir/index.php/jrms/article/view/3978/1852>
- [2] Miller IN, Cronin-Golomb A. Gender differences in Parkinson's disease: Clinical characteristics and cognition. *Movement Disorders* 2010; 15;25(16): 2695-703. [<http://dx.doi.org/10.1002/mds.23388>]
- [3] DeMaagd G, Philip A. Parkinson's disease and its management: part 1: Disease entity, risk factors, pathophysiology, clinical presentation, and diagnosis. *P&T* 2015; 40(8): 504-32. [PMID: 26236139]
- [4] Fereshtehnejad SM, Shafieesabet A, Shafieesabet M, Shahidi GA, Delbari A, Lökk J. Mortality in Iranian patients with Parkinson's Disease: Cumulative impact of cardiovascular comorbidities as one major risk factor. *Parkinson's Disease* 2015; 2015 [<http://dx.doi.org/10.1155/2015/834796>]
- [5] van Nimwegen M, Speelman AD, Hofman-van Rossum EJ, *et al.* Physical inactivity in Parkinson's disease. *J Neurol* 2011; 258(12): 2214-21. [<http://dx.doi.org/10.1007/s00415-011-6097-7>] [PMID: 21614433]
- [6] Roth GA, Johnson C, Abajobir A, *et al.* Global, regional, and national burden of cardiovascular diseases for 10 causes, 1990 to 2015. *J Am Coll Cardiol* 2017; 70(1): 1-25. [<http://dx.doi.org/10.1016/j.jacc.2017.04.052>] [PMID: 28527533]
- [7] Mendis S. Global progress in prevention of cardiovascular disease. *Cardiovasc Diagn Ther* 2017; 7(Suppl. 1): S32-8. [<http://dx.doi.org/10.21037/cdt.2017.03.06>] [PMID: 28529920]
- [8] Yeates K, Lohfeld L, Sleeth J, Morales F, Rajkotia Y, Ogedegbe O. A global perspective on cardiovascular disease in vulnerable populations. *Can J Cardiol* 2015; 31(9): 1081-93. [<http://dx.doi.org/10.1016/j.cjca.2015.06.035>] [PMID: 26321432]
- [9] Press V, Freestone I, George CF. Physical activity: the evidence of benefit in the prevention of coronary heart disease. *QJM* 2003; 96(4): 245-51. [<http://dx.doi.org/10.1093/qjmed/hcg041>] [PMID: 12651969]
- [10] Mora S, Lee IM, Buring JE, Ridker PM. Association of physical activity and body mass index with novel and traditional cardiovascular biomarkers in women. *JAMA* 2006; 295(12): 1412-9. [<http://dx.doi.org/10.1001/jama.295.12.1412>] [PMID: 16551713]
- [11] Gidding SS. Physical activity, physical fitness, and cardiovascular risk factors in childhood. *Am J Lifestyle Med* 2007; 1(6): 499-505. [<http://dx.doi.org/10.1177/1559827607306432>]
- [12] Choo CC, Chew PKH, Lai SM, *et al.* Effect of cardiac rehabilitation on quality of life, depression and anxiety in Asian patients. *Int J Environ Res Public Health* 2018; 15(6): 1095. [<http://dx.doi.org/10.3390/ijerph15061095>] [PMID: 29843421]
- [13] Bocalini DS, dos Santos L, Serra AJ. Physical exercise improves the functional capacity and quality of life in patients with heart failure. *Clinics (São Paulo)* 2008; 63(4): 437-42. [<http://dx.doi.org/10.1590/S1807-59322008000400005>] [PMID: 180759322008000400005]

- 18719752]
- [14] Mogharnasi M, Bagheri M. The effect of 12 weeks of circuit resistance training on C-reactive protein and lipid profile in inactive women. *J Sports Biosci* 2014; 6: 233-44.
- [15] Ellis T, Boudreau JK, DeAngelis TR, *et al.* Barriers to exercise in people with Parkinson disease. *Phys Ther* 2013; 93(5): 628-36. [<http://dx.doi.org/10.2522/ptj.20120279>] [PMID: 23288910]
- [16] Marchese R, Hill A. *The Essential Guide to Fitness for the Fitness Instructor*. Pearson Education 2005.
- [17] Fathi M, Hejazi K. The effect of Eight weeks aerobic training on Resistin levels and cardio respiratory fitness in sedentary middle-aged women. *Med j of Mashhad univ* 2015; 58(9): 489-497.
- [18] Taheri L. The Effects of 8week aerobic exercise on blood lipoprotein of non-athletic middle-aged women of ahvaz. *Harakat* 2007; 9(1): 87-99.
- [19] Soori R, Rezaeian N, Montazeri H. Effects of resistance and endurance training on coronary heart disease biomarker in sedentary obese women. *Majallah-i Ghudad-i Darun/Riz va Mitabulism-i Iran* 2011; 13(2): 179-89.
- [20] Liao F, An R, Pu F, Burns S, Shen S, Jan YK. Effect of exercise on risk factors of diabetic foot ulcers: A systematic review and meta-analysis. *American journal of physical medicine & rehabilitation* 2019; 98(2): 103-6. [<http://dx.doi.org/10.1097/PHM.0000000000001002>]
- [21] Nayeibifar S, Afzalpour ME, Saghebjo M, Hedayati M, Shirzaee P. The effect of aerobic and resistance trainings on serum C-Reactive Protein, lipid profile and body composition in overweight women. *Modern Care J* 2012; 15;8(4): 186-96.
- [22] Hoseini K. Effect of resistance training, aerobic and anaerobic training on lipid profile and CRP in obese girls. *J of Sabzevar Univ Med Sci* 2011; 18(3): 188-97.
- [23] Rahimi N, Marandi SM, Esfarjani F, Ghasemi GA, Habibi N. The effect of one cycle of yoga exercises on blood lipid of the female patients with type II diabetes. *Sadra Med Sci J* 2015; 2(3): 223-4.
- [24] Etemad Z, Esmailnasab N. The effects of strength exercise program of anterior muscles on body composition and serum lipids. *Majallah-i Ilmi-i Danishgah-i Ulum-i Pizishki-i Kurdistan* 2010; 14(4): 20-8.
- [25] Saadatnia A, Ebrahim K, Izadpanah N. Comparing Effect of Aerobic and Resistance Training on Profile of Apolipoprotein and Lipoprotein in Overweight Females. *J Shahid Sadoughi Univ Med Sci* 2015; 23(2): 1887-94.
- [26] Baharloo S, Taghian F, Hedayati M. Effects of aerobic exercise on c-reactive protein and lipid profile in subclinical hypothyroidism among overweight-obese women. *Modares J Med Sci: Pathobiology* 2014; 17(1)
- [27] Askari A, Askari B, Fallah Z. Effect of eight weeks aerobic training on serum lipid and lipoprotein levels in women. *J Gorgan Univ Med Sci* 2012; 14(1): 26-32.
- [28] Maggio AB, Aggoun Y, Martin XE, Marchand LM, Beghetti M, Farpour-Lambert NJ. Long-term follow-up of cardiovascular risk factors after exercise training in obese children. *Int J Pediatr Obes* 2011; 6(2-2): e603-10. [<http://dx.doi.org/10.3109/17477166.2010.530665>] [PMID: 21142752]
- [29] Resaland GK, Anderssen SA, Holme IM, Mamen A, Andersen LB. Effects of a 2-year school-based daily physical activity intervention on cardiovascular disease risk factors: the Sogndal school-intervention study. *Scand J Med Sci Sports* 2011; 21(6): e122-31. [<http://dx.doi.org/10.1111/j.1600-0838.2010.01181.x>] [PMID: 22126720]
- [30] Azizi M, Hoseini R. Relationship between physical activity level and risk factors of cardiovascular disease in male college students. *J Pract Stud of Biosci in Sport* 2013; 1(2): 110-23.
- [31] Rahmani-Nia F, Rahnema N, Hojjati Z, Soltani B. Acute effects of aerobic and resistance exercises on serum leptin and risk factors for coronary heart disease in obese females. *Sport Sci Health* 2008; 2(3): 118. [<http://dx.doi.org/10.1007/s11332-008-0050-x>]
- [32] Abdi Keykanlo N, Rohani H, Asari F. Effects of 8 weeks aerobic training on body composition and plasma levels of insulin-like growth factor-1 and insulin-like growth factor binding protein-3 in obese women. *koomesh* 2014; 15(3): 302-309.
- [33] Attarzadeh Hoeini SR, Hojati Oshrovani Z, Soltani H, Hossein Kakhk SA. Changes in pulmonary function and peak oxygen consumption in response to interval aerobic training in sedentary girls. *Quarterly J Sabzevar Univ Med Sci* 2012; 19(1): 42-51.
- [34] Najaran AF, NourElahi Moghadam H, Mortazian M, Haj Fathali AR, Jamshidi M. Evaluation of Effect of Levodopa on Heart Rate Variability in Parkinson's Disease. *ANN Military Health Sci Res* 2007; 5(3): 1287-92.
- [35] Kazemi T, Rafatpanah N. Management of blood pressure in Parkinson's disease: Need for special attention of a neurologist. *J Birjand Univ Med Sci* 2016; 23(4): 349-52.
- [36] Rahimian MZ, attarzadeh HR, arianezhad J. The effect of aerobic training and diet on cardiovascular risk factors and blood pressure in overweight and obese women with hypertension. *Arch Iran Med* 2011; 14(3)
- [37] Arazi H, Jorbonian A, Asghari E. Comparison of concurrent (resistance-aerobic) and aerobic training on vo2max lipid profile, blood glucose and blood pressure in middle-aged men at risk for cardiovascular disease. *J Shahid Sadoughi Univ Med Sci* 2013; 15;20(5): 627-38.
- [38] Johansen KL. Exercise and chronic kidney disease: Current recommendations. *Sports Med* 2005; 35(6): 485-99. [<http://dx.doi.org/10.2165/00007256-200535060-00003>] [PMID: 15974634]
- [39] Smeltzer S, Bare B, Hinkle J, Cheever K. *China: Lippincott Williams and Wilkins*; 2010. Textbook of medical surgical nursing Brunner and suddarth. 889.
- [40] Myers J, Hadley D, Oswald U, *et al.* Effects of exercise training on heart rate recovery in patients with chronic heart failure. *Am Heart J* 2007; 153(6): 1056-63. [<http://dx.doi.org/10.1016/j.ahj.2007.02.038>] [PMID: 17540210]
- [41] Rafati FM, Taghian F, Pakfetrat M. The effect of 8 week running on treadmills on the amount of blood pressure and excreted albumin protein in patients with chronic kidney. *J Army Univ Med Sci* 2012; 10(2): 118-24.